

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 July 2002 (04.07.2002)

PCT

(10) International Publication Number
WO 02/052021 A1

(51) International Patent Classification⁷: C12N 15/61,
9/90, 15/63, 1/20, C12P 19/02

(74) Agent: LEE, Han-Young; 8th Fl., Seowon Bldg., 1675-
1Seocho-dong, Seocho-gu, Seoul 137-070 (KR).

(21) International Application Number: PCT/KR01/02243

(22) International Filing Date:

22 December 2001 (22.12.2001)

(25) Filing Language:

Korean

(26) Publication Language:

English

(30) Priority Data:

2000/80608 22 December 2000 (22.12.2000) KR
2000/80711 18 December 2001 (18.12.2001) KR

(71) Applicant and

(72) Inventor: PYUN, Yu, Ryang [KR/KR]; 117-14, Yeonhee-
dong, Seodaemun-gu, Seoul 120-114 (KR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KIM, Byoung,
Chan [KR/KR]; 1007-1201 Hoogok-maul, Ilsan-3-dong,
Ilsan-gu, Gyeonggi-do 411-735 (KR). LEE, Han,
Seung [KR/KR]; 791-2137 Mia-1-dong, Gangbuk-gu,
Seoul 142-821 (KR). LEE, Dong, Woo [KR/KR]; 294-1
Hongseon-3-dong, Seodaemun-gu, Seoul 120-847 (KR).
LEE, Yoon, Hee [KR/KR]; 882-24 Bangbae-4-dong,
Seocho-gu, Seoul 137-840 (KR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

- with international search report
- entirely in electronic form (except for this front page) and
available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: THERMOSTABLE L-ARABINOSE ISOMERASE AND PROCESS FOR PREPARING D-TAGATOSE THEREBY

(57) Abstract: The present invention relates to a novel gene coding for L-arabinose isomerase derived from *Thermotoga neapolitana* 5068, a thermostable arabinose isomerase expressed from the said gene, a recombinant expression vector containing the said gene, a microorganism transformed with the said expression vector, a process for preparing thermostable arabinose isomerase from the said transformant and a process for preparing D-tagatose employing the said enzyme. Since the recombinant arabinose isomerase of the invention is highly thermostable and can produce tagatose with high yield at high temperature, it can be efficiently applied in pharmaceutical and food industries.

WO 02/052021 A1

**THERMOSTABLE L-ARABINOSE ISOMERASE AND PROCESS FOR
PREPARING D-TAGATOSE THEREBY**

5 BACKGROUND OF THE INVENTION

Field of the Invention

 The present invention relates to a thermostable
10 arabinose isomerase and a process for preparing tagatose
 using the same, more specifically, to a noble gene coding
 for L-arabinose isomerase derived from *Thermotoga*
 neapolitana 5068, a thermostable arabinose isomerase
 expressed from the said gene, a recombinant expression
15 vector containing the said gene, a microorganism
 transformed with the said expression vector, a process for
 preparing thermostable arabinose isomerase from the said
 transformant and a process for preparing D-tagatose
 employing the said enzyme.

20

Background of the Invention

 In recent years, growing concerns about health have
 led much research effort to the development of healthful
25 foods. As one of the above efforts, sugar alcohols have
 been proposed as sweeteners which can substitute sugar,
 known to cause adult diseases, and are practically being
 used. Since the said sweeteners are known to have adverse
 side effects such as causing diarrhea when ingested more
30 than certain amount, there is an urgent need to develop
 substitutional sweeteners without harmful effects.

 Among substitutional sweeteners which have little
 side effect, tagatose, a keto-sugar of galactose, has
 similar sweetness to D-fructose, and has known not to be
35 absorbed or metabolized in the body, making tagatose a safe
 low-caloric substitutional sweetener for sugar. Also, it
 has been reported that tagatose can be employed as an

intermediate for the preparation of useful optically active isomers, detergents and cosmetics, also, as an additive or raw material for the synthesis of drugs, especially, its ability to lower blood sugar level renders tagatose a therapeutic and preventive agent for diabetes, and a low caloric diet agent.

Currently, tagatose is mostly prepared via chemical synthesis from galactose(see: USP 5,002,612), which comprises the steps of isomerization of galactose catalyzed by metal hydroxide in the presence of inorganic salts to form an intermediate of metal hydroxide-tagatose complex, and neutralization of the complex by adding acid to yield final product, tagatose.

Alternative method for manufacturing tagatose is an enzymatic method in which galactose is converted into tagatose via conversion of aldose or aldose derivatives into ketose or ketose derivatives. Especially, it has been reported that arabinose isomerase which catalyzes the conversion reaction of L-arabinose into L-ribulose can be employed for production of tagatose in vitro using galactose as a substrate. However, the yield of tagatose produced by arabinose isomerase from galactose is as low as 20%, hindering industrial application of conversion process of galactose into tagarose. Although the method for manufacturing tagatose from milk or cheese has been developed(see: USP 6,057,135), again, low yield is the limitation for its industrial use.

Under the circumstances, there are strong reasons for exploring and developing a novel enzyme which can produce tagatose with high yield.

Summary of the Invention

The present inventors have made an effort to develop an enzyme which can produce tagatose with high yield, thus, have found that tagatose can be produced with high yield from galactose by employing a recombinant arabinose

isomerase produced from *E. coli* transformed with recombinant vector containing arabinose isomerase gene derived from *Thermotoga neapolitana* 5068.

5 The first object of the present invention is, therefore, to provide arabinose isomerase gene derived from *Thermotoga neapolitana* 5068.

 The second object of the invention is to provide arabinose isomerase expressed from the gene.

10 The third object of the invention is to provide a recombinant expression vector containing the arabinose isomerase gene.

 The fourth object of the invention is to provide a recombinant *E. coli* transformed with the recombinant
15 expression vector.

 The fifth object of the invention is to provide a process for preparing recombinant arabinose isomerase using the transformed *E. coli*.

 The sixth object of the invention is to provide a
20 process for preparing tagatose from galactose using the enzyme.

BRIEF DESCRIPTION OF THE DRAWINGS

25 The above and the other objects and features of the present invention will become apparent from the following descriptions given in conjunction with the accompanying drawings, in which:

30 Figure 1 is a schematic diagram showing the construction strategy of an expression vector containing arabinose isomerase gene of the invention.

 Figure 2 is a graph showing activity profile of arabinose isomerase of the invention depending on
35 temperature.

 Figure 3 is a graph showing thermostability of arabinose isomerase of the invention.

Figure 4 is a graph showing the time course of conversion rate of galactose into tagatose by arabinose isomerase of the invention at various reaction temperatures.

5 Figure 5 is a graph showing the time course of changes in thermostability of immobilized arabinose isomerase of the invention.

DETAILED DESCRIPTION OF THE INVENTION

10

To prepare thermophilic or thermostable arabinose isomerase for industrial use, the present inventors have cloned a gene coding for arabinose isomerase from genomic DNA of *Thermotoga neapolitana* 5068(DSM 5608) and analyzed
15 nucleotide sequence and deduced amino acid sequence from the said gene. The nucleotide sequence and deduced amino acid sequence of the gene encoding arabinose isomerase of the invention(SEQ ID NO: 3) has shown to have 83.2% and
20 94.8% homology, respectively, to those of the putative arabinose isomerase gene of *Thermotoga maritima* of which entire nucleotide sequence has been verified via genome project.

For high level expression of the said cloned arabinose isomerase in *E. coli*, the gene coding for the
25 enzyme was inserted into an expression vector pET22b(+) (Novagen, U.S.A.) to construct a recombinant expression vector pTNAI, which was then introduced into *E. coli* BL21. The transformed recombinant *E. coli* was named "*E. coli* BL21/DE3(pTNAI)" and deposited with an
30 international depository authority, the Korean Culture Center of Microorganisms(KCCM, #361-221 Hongje-1-dong, Seodaemun-gu, Seoul, Republic of Korea) on December 4, 2000 as accession no. KCCM-10231.

The said *E. coli* BL21/DE3(pTNAI) was grown to obtain
35 recombinant arabinose isomerase, which was characterized to have optimum pH of 7.0, optimum reaction temperature of 85°C. Furthermore, over 80% of remaining activity was

measured after 2 hour heat treatment at 80°C, indicating that the enzyme is exceedingly heat stable.

Tagatose can be produced by employing arabinose isomerase of the invention prepared from *E. coli* transformed with a recombinant expression vector containing the gene for arabinose isomerase derived from *Thermotoga* sp., and galactose as a substrate, under a condition of pH 5 to 8, more preferably pH 6 to 8, most preferably pH 7, and 60 to 100°C, more preferably 70 to 95°C, most preferably 85°C.

Aqueous solution of galactose was subjected to isomerization reaction employing recombinant arabinose isomerase of the invention, and it has been found that conversion rate into tagatose was over 68% at 80°C.

When the said recombinant arabinose isomerase is employed for industrial production of tagatose, soluble form of the enzyme may be employed, nevertheless, it is more preferable to immobilize the enzyme on the beads used in industry. For example, in case of the recombinant arabinose isomerase of the invention immobilized on silica beads, the remaining activity was measured to be over 80% of original activity after 20 day-heat treatment at 90°C, thus, it can be applied for thermal process over 80°C in industry.

The present invention is further illustrated in the following examples, which should not be taken to limit the scope of the invention.

Example 1: Cloning of arabinose isomerase gene

Thermotoga neapolitana 5068 (DSM 5068) was grown under an anaerobic condition and cells were harvested by centrifugation at 8000xg for 10 minutes. Genomic DNA isolated from the cells harvested above was partial digested with Sau3AI (TaKaRa Biotechnology, Japan) to obtain 12kb or shorter fragments of DNA. The DNA fragments

were inserted into ZAP Expression Vector(Stratagene, U.S.A.) and packaged to prepare a genomic library of *Thermotoga neapolitana* 5068. Nucleotide sequences of the genes for conventional thermophilic or thermostable arabinose isomerase were analyzed to prepare primer araAF: 5'-ATGATCGATCTCAAACAGTATGAG-3'(SEQ ID NO: 1) and primer araAR: 5'-TCATCTTTTAAAGTCCCC-3'(SEQ ID NO: 2), which were used in PCR for the preparation of probes for DNA-DNA hybridization. The genomic library prepared above was screened for DNA fragments containing arabinose isomerase gene by DNA-DNA hybridization to obtain a recombinant vector containing a gene encoding arabinose isomerase of *Thermotoga neapolitana* 5068. The nucleotide sequence of arabinose isomerase gene(SEQ ID No: 3) cloned above and the deduced amino acid sequence(SEQ ID No: 4) from the said gene were compared with those of known arabinose isomerase genes, respectively(see: Table 1).

Table 1: Comparison of homology between arabinose isomerase of the invention and known arabinose isomerases

Strain	Gene Sequence (homology, %)	Amino Acid Sequence (homology, %)
<i>Thermotoga maritima</i>	83.2	94.8
<i>Bacillus stearothermophilus</i>	61.9	62.8
<i>Bacillus halodurans</i>	59.1	59.0
<i>Bacillus subtilis</i>	58.6	55.5
<i>Salmonella typhimurium</i>	57.8	54.5
<i>Escherichia coli</i>	59.0	54.3
<i>Mycobacterium smegmatis</i>	56.3	50.7

As shown in Table 1, it has been found that the arabinose isomerase of the invention is a novel enzyme which has 83.2% homology of nucleotide sequence and 94.8%

homology of amino acid sequence to the sequences of published putative arabinose isomerase of *Thermotoga maritima*, respectively.

5 Example 2: Preparation of recombinant expression vector and recombinant *E. coli*

In order to obtain high level expression of the said thermostable arabinose isomerase in *E. coli* using the
10 arabinose isomerase gene obtained in Example 1, the said gene was inserted into an expression vector pET 22b(+) (Novagen, U.S.A.) double-digested with *NdeI* and *EcoRI* to construct a recombinant expression vector pTNAI (see: Figure 1), which was then introduced into *E.*
15 *coli* BL21. The transformed recombinant *E. coli* was named "*E. coli* BL21/DE3(pTNAI)" and deposited with an international depository authority, the Korean Culture Center of Microorganisms (KCCM, #361-221 Hongje-1-dong, Seodaemun-gu, Seoul, Republic of Korea) on December 4,
20 2000 as accession no. KCCM-10231.

Example 3: Expression of recombinant arabinose isomerase

The recombinant *E. coli* BL21/DE3(pTNAI) (KCCM-10231)
25 prepared in Example 2 was inoculated into LB (Luria-Bertani) medium (1% v/v) and incubated at 37°C for 2 hours, to which lactose was added to a final concentration of 1mM and expression of recombinant arabinose isomerase was induced for 12 hours. For assay of expressed arabinose
30 isomerase, cells were collected by centrifugation at 8000xg for 10 minutes, resuspended in 10ml of 100mM MOPS buffer (4-morpholinepropanesulfonic acid, pH 7.0), and then disrupted by sonication to obtain crude enzyme, with which galactose isomerization reaction was carried
35 out. Galactose isomerization was performed by mixing 100μl of the said crude enzyme solution with 40mM (final concentration) galactose as a substrate, followed by

adding 1ml of enzyme reaction buffer(100mM MOPS buffer, pH 7.0) containing cofactors(1mM MnCl_2 , 1mM CoCl_2) and incubating at 85°C for 20 minutes. The product of the above reaction was detected using cysteine-carbazole-sulfuric acid method(see: Dische, Z., and E. Borenfreund., A New Spectrophotometric Method for the Detection and Determination of Keto Sugars and Trioses, *J. Biol. Chem.*, 192:583-587, 1951), and it has been found that normal galactose isomerization has been undergone.

10

Example 4: Purification of recombinant arabinose isomerase

For purification of recombinant arabinose isomerase expressed by the method described in Example 3, cells were collected by centrifugation at 8000xg for 19 minutes and cell wall of *E. coli* was disrupted by sonication, which was followed by centrifugation at 20,000xg for 20 minutes to obtain supernatant. Then, the said supernatant was heat-treated at 85°C for 20 minutes, centrifuged at 20,000xg for 20 minutes to get rid of precipitate, and the supernatant was further purified by ammonium sulfate-mediated precipitation and finally ion-exchange column chromatography(Q-Sepharose Fast Flow, Pharmacia, Sweden). pH dependancy of the said purified enzyme was analyzed and optimum pH was found to be around 7.0.

Example 5: Optimum pH and optimum temperature of recombinant arabinose isomerase

Activity of the purified recombinant arabinose isomerase prepared in Example 4 was analyzed on galactose substrate and optimum pH was found to be around 7.0. Optimum temperature for isomerization reaction was determined using the same method as described in Example 3. The tested reaction temperatures for galactose isomerization were 60, 70, 75, 80, 85, 90 and 100°C, and maximum activity was obtained around 85°C(see: Figure 2).

Example 6: Thermostability of recombinant arabinose
isomerase

5 To assess the thermostability of recombinant
arabinose isomerase of the invention, crude enzyme
prepared in Example 3 was heat-treated at 60, 70, 80 and
90°C for 10, 20, 30, 60, 90 and 120 minutes respectively,
and remaining activity of recombinant arabinose isomerase
10 for isomerization was determined as described in Example
3(see: Figure 3). As shown in Figure 3, it has been found
that over 80% of enzyme activity was remained after 2 hour
heat-treatment at 80°C.

15 Example 7: Conversion rate of galactose into tagatose at
various temperature

By employing recombinant arabinose isomerase of the
invention, the conversion rate of galactose into tagatose
20 was determined at various temperatures and various time
points. Substrate used was 10mM galactose instead of 40mM
galactose in enzyme reaction mixture in Example 3. After
incubation at 60, 70, 80 and 90°C for 20 hours, tagatose
yield was determined employing BioLC(see: Table 2 and
25 Figure 4).

Table 2: Conversion rate of galactose into tagatose at
various temperature

Enzyme Reaction Temperature	60°C	70°C	80°C	90°C
Conversion Rate into Tagatose	31.7	40.4	68.1	57.4

30

As shown in Table 2 and Figure 4, the higher the
reaction temperature was, the higher tagatose yield was
obtained, and conversion rate into tagatose was as high as
68% at 80°C.

35

Example 8: Immobilization of arabinose isomerase and improvement of thermostability

Arabinose isomerase was immobilized on silica beads,
5 heat-treated under an aqueous condition at 90°C and the
remaining activity was determined at various time
points(see: Figure 5). As shown in Figure 5, remaining
activity of the immobilized enzyme was over 80% after 20
day-heat treatment at 90°C and over 60% after 30 day-heat
10 treatment, indicating that the immobilized arabinose
isomerase of the invention can be applied for thermal
process in industry.

As clearly illustrated and demonstrated above, the
15 present invention provides a noble gene coding for L-
arabinose isomerase derived from *Thermotoga neapolitana* 5068,
a thermostable arabinose isomerase expressed from the said
gene, a recombinant expression vector containing the said
gene, a microorganism transformed with the said expression
20 vector, a process for preparing thermostable arabinose
isomerase from the said transformant and a process for
preparing D-tagatose employing the said enzyme. Since the
recombinant arabinose isomerase of the invention is highly
thermostable and can produce tagatose with high yield at
25 high temperature, it can be efficiently applied in
pharmaceutical and food industries.

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL
(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in description On page 4 , lines 23-33 and page 7 , lines 8-20 .	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on additional sheet	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution (including postal code and country) Korean Culture Center of Microorganisms(KCCM) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu Seoul, 120-091, Republic of Korea	
Date of deposit Dec. 04, 2000	Accession Number KCCM-10231
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information continues on an additional sheet	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

5

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

What is Claimed is:

1. A gene coding for arabinose isomerase having a nucleotide sequence of SEQ ID NO: 3 which is derived from
5 *Thermotoga* sp.

2. A gene coding for arabinose isomerase having a nucleotide sequence of SEQ ID NO: 4 which is derived from the nucleotide sequence of claim 1.
10

3. A recombinant expression vector pTNAI represented as a genetic map of Fig. 1 which contains the nucleotide sequence of SEQ ID NO: 3 of a gene derived from *Thermotoga* sp.
15

4. *E. coli* BL21/DE3(pTNAI) (KCCM-10231) transformed with the recombinant expression vector pTNAI of claim 3.

5. A process for preparing a recombinant arabinose isomerase which comprises a step of culturing a microorganism transformed with a recombinant expression vector containing the gene for arabinose isomerase derived from *Thermotoga* sp. of claim 1 to obtain a recombinant arabinose isomerase from the culture.
20

6. The process for preparing a recombinant arabinose isomerase of claim 5, wherein the microorganism transformed with a recombinant expression vector containing the gene for arabinose isomerase is *E. coli* BL21/DE3(pTNAI) (KCCM-
25 10231).
30

7. A process for preparing tagatose which comprises a step of reacting arabinose isomerase prepared from a microorganism transformed with a recombinant expression vector containing a gene for arabinose isomerase derived from *Thermotoga* sp. with a substrate of galactose under a condition of pH 5 to 8 and 50 to 100°C to obtain tagatose.
35

8. The process for preparing tagatose of claim 7,
wherein the microorganism transformed with a recombinant
expression vector containing a gene for arabinose isomerase
5 derived from *Thermotoga* sp. is *E. coli* BL21/DE3(pTNAI)
(KCCM-10231).

9. The process for preparing tagatose of claim 7,
wherein the arabinose isomerase is immobilized recombinant
10 arabinose isomerase.

15

20

25

30

35

1/3




Fig. 1

2/3




Fig. 2




Fig. 3

3/3




Fig. 4




Fig. 5

SEQUENCE LISTING

<110> PYUN, YU RYANG, et al.

5 <120> Thermostable L-Arabinose Isomerase and Process for Preparing D-ta
gatose thereby

<150> KR 10-2000-0080608

10 <151> 2000-12-22

<150> KR 10-2001-0080711

<151> 2001-12-18

15 <160> 4

<170> KopatentIn 1.6

<210> 1

20 <211> 24

<212> DNA

<213> Artificial Sequence

<220>

25 <223> primer araAF

<400> 1

atgatcgatc tcaaacagta tgag 24

30 <210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

35 <220>

<223> primer araAR

	<400>	2	
	tcacatcttttt	aaaagtcctcc	20
5	<210>	3	
	<211>	1491	
	<212>	DNA	
	<213>	Thermotoga neapolitana 5068	
10	<400>	3	
	atgatcgatc	tcaaacagta tgagttctgg tttcttgtcg gcagccagta tctctacggt	60
	ctggagacgt	tgaagaaggt agagcagcag gcaagcagga tagttgaggc actgaacaat	120
15	gatcccatct	ttccctcaaa gatcgttctg aaacctgttc tgaaaaattc cgccgagatc	180
	agagagatct	tcgaaaaggc aaatgcagaa ccaaaatgcg ccggtgtcat cgtgtggatg	240
20	cacacgttct	caccttcgaa gatgtggata agaggcctct ccatcaataa aaaacccttg	300
	cttcacctcc	acaccagta caacagggag atcccgtggg acacgatcga tatggactac	360
	atgaacctga	accaatctgc ccacggtgac agggaaacacg gattcattca cgcgaggatg	420
25	agactcccaa	gaaaggtcgt ggtgggacat tgggaagaca gagaagtcag ggaaaagatc	480
	gcaaaatgga	tgagagtggc ctgcgcgata caggatggaa gaactggaca gatcgtgaga	540
30	ttcggcgata	acatgagaga ggttgccagc accgaagacg acaagggtga ggcacagata	600
	aaactcggct	ggtccataaa cacctgggggt gtcggagagc tcgccgaggg agtgaaggcg	660
	gttcagaaa	acgaagtgga ggaattgttg aaggagtaca aagaaaggta catcatgcca	720
35	gaagacgaat	acagcctcaa agcgatcaga gaacaggcga agatggagat tgcactgaga	780

	gagtttctga aagagaagaa tgccatcgcc ttcaccacca ccttcgagga tcttcacgat	840
	cttccccagc ttcccgtct tgcagtcagc aggcctcatgg aggaagggtg tggatttgga	900
5	gcggaaggag actggaaggc agccgggctt gtgagggtt tgaaggcat gggagctggt	960
	cttcccgtg gtacatcctt catggaggac tacacctacc atctcacacc gggaaacgaa	1020
	ctcgtgctgg gagcgacat gctagagggtg tgccccacga tcgctaagga aaagccaaga	1080
10	atagagggtgc atcctctcag catcggtgga aaagcagatc ctgcacgcct tgttttcgat	1140
	ggacaagaag gtcccgtgt caacgcctcc atcgttgaca tgggaaacag gttcaggctg	1200
15	gtagtgaaca gagtgttgtc tgttccatt gaaaggaaga tgcccaaact tccaacggca	1260
	agagttttgt ggaagccgct tctgatttc aagagggtga cgactgcgtg gattctcgt	1320
	ggaggatccc atcatactgc cttctcaaca gcggtggatg tggagtacct catcgactgg	1380
20	gcggaggctt tggagataga gtatcttgtc atcgatgaaa atctggatct ggagaacttc	1440
	aaaaaggaac tgagatggaa cgaactctac tggggacttt taaaaagatg a	1491
25	<div> <div><210> 4</div> <div><211> 496</div> <div><212> PRT</div> <div><213> Thermotoga neapolitana 5068</div> </div>	
30	<div> <div><400> 4</div> <div>Met Ile Asp Leu Lys Gln Tyr Glu Phe Trp Phe Leu Val Gly Ser Gln</div> <div>1 5 10 15</div> </div>	
35	<div> <div>Tyr Leu Tyr Gly Leu Glu Thr Leu Lys Lys Val Glu Gln Gln Ala Ser</div> <div>20 25 30</div> </div>	

Arg Ile Val Glu Ala Leu Asn Asn Asp Pro Ile Phe Pro Ser Lys Ile
35 40 45

Val Leu Lys Pro Val Leu Lys Asn Ser Ala Glu Ile Arg Glu Ile Phe
5 50 55 60

Glu Lys Ala Asn Ala Glu Pro Lys Cys Ala Gly Val Ile Val Trp Met
65 70 75 80

10 His Thr Phe Ser Pro Ser Lys Met Trp Ile Arg Gly Leu Ser Ile Asn
85 90 95

Lys Lys Pro Leu Leu His Leu His Thr Gln Tyr Asn Arg Glu Ile Pro
100 105 110

15 Trp Asp Thr Ile Asp Met Asp Tyr Met Asn Leu Asn Gln Ser Ala His
115 120 125

Gly Asp Arg Glu His Gly Phe Ile His Ala Arg Met Arg Leu Pro Arg
20 130 135 140

Lys Val Val Val Gly His Trp Glu Asp Arg Glu Val Arg Glu Lys Ile
145 150 155 160

25 Ala Lys Trp Met Arg Val Ala Cys Ala Ile Gln Asp Gly Arg Thr Gly
165 170 175

Gln Ile Val Arg Phe Gly Asp Asn Met Arg Glu Val Ala Ser Thr Glu
180 185 190

30 Asp Asp Lys Val Glu Ala Gln Ile Lys Leu Gly Trp Ser Ile Asn Thr
195 200 205

Trp Gly Val Gly Glu Leu Ala Glu Gly Val Lys Ala Val Pro Glu Asn
35 210 215 220

Glu Val Glu Glu Leu Leu Lys Glu Tyr Lys Glu Arg Tyr Ile Met Pro

225 230 235 240
Glu Asp Glu Tyr Ser Leu Lys Ala Ile Arg Glu Gln Ala Lys Met Glu
 245 250 255
5
Ile Ala Leu Arg Glu Phe Leu Lys Glu Lys Asn Ala Ile Ala Phe Thr
 260 265 270
Thr Thr Phe Glu Asp Leu His Asp Leu Pro Gln Leu Pro Gly Leu Ala
10 275 280 285
Val Gln Arg Leu Met Glu Glu Gly Tyr Gly Phe Gly Ala Glu Gly Asp
 290 295 300
15 Trp Lys Ala Ala Gly Leu Val Arg Ala Leu Lys Val Met Gly Ala Gly
 305 310 315 320
Leu Pro Gly Gly Thr Ser Phe Met Glu Asp Tyr Thr Tyr His Leu Thr
 325 330 335
20
Pro Gly Asn Glu Leu Val Leu Gly Ala His Met Leu Glu Val Cys Pro
 340 345 350
Thr Ile Ala Lys Glu Lys Pro Arg Ile Glu Val His Pro Leu Ser Ile
25 355 360 365
Gly Gly Lys Ala Asp Pro Ala Arg Leu Val Phe Asp Gly Gln Glu Gly
 370 375 380
30 Pro Ala Val Asn Ala Ser Ile Val Asp Met Gly Asn Arg Phe Arg Leu
 385 390 395 400
Val Val Asn Arg Val Leu Ser Val Pro Ile Glu Arg Lys Met Pro Lys
 405 410 415
35
Leu Pro Thr Ala Arg Val Leu Trp Lys Pro Leu Pro Asp Phe Lys Arg
 420 425 430

Ala Thr Thr Ala Trp Ile Leu Ala Gly Gly Ser His His Thr Ala Phe
435 440 445

5 Ser Thr Ala Val Asp Val Glu Tyr Leu Ile Asp Trp Ala Glu Ala Leu
450 455 460

Glu Ile Glu Tyr Leu Val Ile Asp Glu Asn Leu Asp Leu Glu Asn Phe
465 470 475 480

10 Lys Lys Glu Leu Arg Trp Asn Glu Leu Tyr Trp Gly Leu Leu Lys Arg
485 490 495

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR01/02243

A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C12N 15/61, C12N 9/90, C12N 15/63, C12N 1/20, C12P 19/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N 15/61, C12N 9/90, C12N 15/63, C12N 1/20, C12P 19/02

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

NCBI, PubMed, CA, USPTO, PAJ, Espacenet, "Thermotoga", "arabinose isomerase", "tagatose"

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A Y	Nelson, K.E. et al., "Evidence for lateral gene transfer between Archaea and bacteria from genome sequence of <i>Thermotoga maritima</i> ", Nature, 399(6734), 323-29, 1999 & NCBI Accession # AE001709.	1-6, 8, 9 7
Y	Roh, H.J. et al., "Bioconversion of D-galactose into D-tagatose by expression of L-arabinose isomerase", Biotechnol. Appl. Biochem., 31(1), 1-4, 2000.	7
Y	US 6,057,135 A (Kraft-Foods, Inc.), 02 May 2000.	7
Y	WO 2000/068397 A1 (Tongyang Confectionery Co.), 16 Nov. 2000.	7

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 MARCH 2002 (13.03.2002)

Date of mailing of the international search report

13 MARCH 2002 (13.03.2002)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
Government Complex-Daejeon, 920 Dunsan-dong, Seo-gu,
Daejeon Metropolitan City 302-701, Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

LEE, Cheo Young

Telephone No. 82-42-481-5594



Information on patent family members

PCT/KR01/02243

Form PCT/ISA/210 (patent family annex) (July 1998)